

Clinical Policy: Mogamulizumab-kpkc (Poteligeo)

Reference Number: PA.CP.PHAR.139

Effective Date: 10/2018

Last Review Date: 10/2024

Description

Mogamulizumab-kpkc (Poteligeo[®]) is a CC chemokine receptor type 4 (CCR4)-directed monoclonal antibody.

FDA Approved Indication(s)

Poteligeo is indicated for the treatment of adult patients with relapsed or refractory mycosis fungoides (MF) or Sézary syndrome (SS) after at least one prior systemic therapy.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of PA Health & Wellness[®] that Poteligeo is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Mycosis Fungoides/Sézary Syndrome (must meet all):

1. Diagnosis of MF or SS;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. Request meets one of the following (a or b):
 - a. Dose does not exceed 1 mg/kg on days 1, 8, 15, and 22 of the first 28-day cycle and on days 1 and 15 of each subsequent cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 6 months

B. Adult T-Cell Leukemia/Lymphoma (off-label) (must meet all):

1. Diagnosis of adult T-cell leukemia/lymphoma (ATLL);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. Prescribed as a single-agent;
5. Failure of a first-line chemotherapy regimen (*see Appendix B for examples*);
**Prior authorization may be required.*
6. Request meets one of the following (a or b):
 - a. Dose does not exceed 1 mg/kg on days 1, 8, 15, and 22 of the first 28-day cycle and on days 1 and 15 of each subsequent cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 6 months

C. Other diagnoses/indications

1. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.PHARM.01) applies;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):
 - a. New dose does not exceed 1 mg/kg on days 1 and 15 of each 28-day cycle;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via PA Health & Wellness benefit or member has previously met all initial approval criteria or the Continuity of Care Policy (PA.PHARM.01) applies.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): PA.CP.PMN.53.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – PA.CP.PMN.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ATLL: adult T-cell leukemia/lymphoma	MF: mycosis fungoides
CCR4: CC chemokine receptor type 4	NCCN: National Comprehensive Cancer Network
CTCL: cutaneous T-cell lymphoma	SS: Sézary syndrome
FDA: Food and Drug Administration	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
ATLL: examples of first-line therapy: <ul style="list-style-type: none"> • Brentuximab vedotin + CHP (cyclophosphamide, doxorubicin, and prednisone) 	Varies	Varies

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<ul style="list-style-type: none"> • CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) • CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, prednisone) • Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) • HyperCVAD (cyclophosphamide, vincristine, doxorubicin, dexamethasone) alternating with high-dose methotrexate and cytarabine • Zidovudine and interferon 		

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
MF, SS	1 mg/kg IV over at least 60 minutes on days 1, 8, 15, and 22 of the first 28-day cycle and on days 1 and 15 of each subsequent cycle until disease progression or unacceptable toxicity	1 mg/kg/dose

VI. Product Availability

Solution for injection in a single-dose vial: 20 mg/5 mL (4 mg/mL)

VII. References

1. Poteligeo Prescribing Information. Bedminster, NJ: Kyowa Kirin, Inc.; March 2023. Available at: <https://www.poteligeohcp.com>. Accessed July 15, 2024.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed August 20, 2024.
3. National Comprehensive Cancer Network. Primary Cutaneous Lymphomas Version 2.2024. Available at: https://www.nccn.org/professionals/physician_gls/pdf/primary_cutaneous.pdf. Accessed August 20, 2024.
4. National Comprehensive Cancer Network. T-Cell Lymphomas Version 4.2024. Available at: https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf. Accessed August 20, 2024.

Reviews, Revisions, and Approvals	Date
Policy created	10/2018
4Q 2019 annual review: No changes per Statewide PDL implementation 01-01-2020	10/2019
4Q 2020 annual review: added age limit; references reviewed and updated.	10/2020

Reviews, Revisions, and Approvals	Date
4Q 2021 annual review: no significant changes; references reviewed and updated.	10/2021
4Q 2022 annual review: no significant changes; references reviewed and updated.	10/2022
4Q 2023 annual review: no significant changes; references reviewed and updated.	10/2023
4Q 2024 annual review: for ATLL initial criteria, added “prescribed as a single-agent” to align with NCCN compendium and guideline; for Appendix B, added another therapeutic option “Zidovudine and interferon” for ATLL first line therapy; references reviewed and updated.	10/2024